


JPL PUBLICATION 82-44, VOLUME IV

Planetary Geometry Handbook

Jupiter Positional Data, 1985-2020

Andrey B. Sergeyevsky
Gerald C. Snyder
Barbara L. Paulson
Ross A. Cunniff



October 1, 1983

NASA

National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

The research described in this publication was carried out by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Abstract

This document contains graphical data necessary for the analysis of planetary exploration missions to Jupiter. Positional and geometric information spanning the time period from 1985 through 2020 is presented; the text explains the data and its usage. This volume is one in a planned series covering planetary mission targets.

Preface

This publication is one in a series of volumes devoted to planetary positional and geometric data. The present Volume IV provides information describing the planet Jupiter. Other volumes to be published in 1983 are II, III, and V, describing Venus, Mars, and Saturn geometric data, respectively. The presentations of planetary positional and geometric data for Mercury (Volume I), Uranus (Volume VI), Neptune (Volume VII), and the Pluto/Charon System (Volume VIII) will be published later.

Contents

I. Introduction	1
II. Description of the Data	1
A. General Comments	1
B. Data Presentation	1
C. Definition of Terms	1
D. Description of Individual Plots	4
III. Application of the Data Presented	6
A. Geocentric Declination of Target Body	6
B. Planetocentric Declination of Earth and Sun	7
C. Right Ascension of Target Planet, Sun, and Earth	7
D. Sun- or Earth-to-Planet Distance	7
E. Heliocentric Longitude	7
F. Angles Between Sun, Earth, and Planet (SEP and ESP)	7
G. Cone and Clock Angles	8
H. Station Rise and Set	8
IV. Astrodynamic Constants	8
Acknowledgments	10
References	11
Figures	
1. Definition of vernal equinox	2
2. Definition of Sun- and Earth-to-planet distance, and Earth-Sun- and Sun-Earth-planet angles at any epoch	3
3. Definition of cone and clock angles	3
4. Schematic of a distant object's rise and set geometry, with a mask; polar view	4
5. Circumpolar and inaccessible regions of the sky for a station observer; meridional view	5
6. Definition of opposition and conjunction events involving Sun, Earth, and planet	8

7. Definition of the mean elements of a planetary orbit	9
8. IAU definition of planetary pole and prime meridian at time d_{50} ..	10

Tables

1. Deep Space Network tracking stations	5
2. Non-DSN tracking stations	5

Positional Data

Jupiter

1985–2020	13
1985	15
1986	21
1987	27
1988	33
1989	39
1990	45
1991	51
1992	57
1993	63
1994	69
1995	75
1996	81
1997	87
1998	93
1999	99
2000	105
2001	111
2002	117
2003	123
2004	129
2005	135
2006	141
2007	147
2008	153
2009	159

2010	165
2011	171
2012	177
2013	183
2014	189
2015	195
2016	201
2017	207
2018	213
2019	219
2020	225

Positional Data

Earth

1985–2020	231
------------------	-----

I. Introduction

The purpose of this series of planetary geometry handbooks is to provide mission and science planners, as well as trajectory designers, with graphical information sufficient for preliminary mission design and evaluation. It is intended to be used in conjunction with the appropriate volumes and parts of the Mission Design Handbooks, Ref. 1, which describe trajectory aspects of the particular missions.

In most respects, this planetary geometry series represents a continuation of the second parts of each of the three volumes of Ref. 2. It extends their coverage by commencing in 1985 and continuing to the year 2020. This time span was chosen to provide sufficient mission duration flexibility for all Earth departures through 2005, as presented in Ref. 1.

The series will consist of eight volumes, each describing a planet as follows:

Volume	Target planet
I	Mercury
II	Venus
III	Mars
IV	Jupiter
V	Saturn
VI	Uranus
VII	Neptune
VIII	Pluto/Charon System

The present Volume IV is devoted to data on the positions of the planet Jupiter. It presents information characterizing the geocentric and heliocentric positional geometry of a spacecraft while it is in the vicinity of the subject planetary body. The spacecraft could encompass a lander, an entry probe, or a flyby vehicle at encounter. Such wide usage of the data is possible because planetary positions change but slowly, while the distances to Earth or Sun are always large compared to those between the spacecraft and the target planet during the encounter.

II. Description of the Data

A. General Comments

All plots for this handbook series were computer generated, to reduce production cost. A program originally developed for Mariner 10 (MVM) and later used to produce the positional

plots for Ref. 2 has been used, with some modifications, on this series as well.

The major change in the new version of the program was the incorporation of the numerically integrated JPL Planetary DE-118 Ephemeris (Ref. 3) as the source for positional planetary information. An analytical and thus less-precise ephemeris, based on the mean elements of Ref. 4, was used in the preceding set of volumes, Ref. 2.

B. Data Presentation

The data in this volume are grouped into 36 subsections, each covering a specific calendar year. The data commence each year on January 1.0, 0^h Greenwich mean time (GMT) and continue through the year until "Day 380," thus providing a two-week overlap into the next year. There are eight plots presented for each year, labeled as follows:

- (1) *Declination* (DECLIN).
- (2) *Right ascension* (RT.ASC).
- (3) *Distance* (DISTANCE).
- (4) *Ecliptic longitude* (EC.LON).
- (5) *Sun-Earth-planet angles* (SUN-EARTH-PLANET).
- (6) *Cone and clock angles* (CA and KA, respectively).
- (7) *Station rise/set times* (STATION RISE/SET) for the three DSN stations (GMT, HRS).
- (8) Same as (7) for three other stations (Weilheim, Nobeyama, and Arecibo).

Twenty-six variables, relating positions of Earth, Sun, target planet (Jupiter), the star Canopus, and six tracking stations, are depicted on these plots for each subsection.

C. Definition of Terms

Planetary positional geometric data can be represented by directions to specified bodies and by angles between some of these directions, both shown as functions of calendar time. To distinguish among these bodies, the following subscripts are introduced:

- (1) E = Earth.
- (2) S = Sun.
- (3) P = planet (Jupiter in this volume).

Two fundamental planes are used in astrodynamics: that of the Earth's orbit about the Sun and that of the Earth's equator. The same two geometric concepts can be extended to all celestial bodies.

The plane of the mean Earth's orbit (EMO) is also called the ecliptic; the term "mean" refers to an averaging process, which eliminates minor short-period oscillations of the orbit pole; these oscillations are caused by repetitive perturbing accelerations caused by solar system objects other than the Sun.

The rotational, angular-momentum vector of each planet defines the polar-axis orientation of that celestial body, but not necessarily its north pole. The plane normal to this axis of rotation is defined as the planetary equator of that body. The latest (1983) International Astronomical Union (IAU) definition of planetary poles is such that the *north pole* of a body is that pole that points north of the Invariable (or Laplacian) Plane, defined as the plane normal to the total angular momentum vector of the solar system. This plane is very close to the ecliptic, and deviates from the latter by less than 0.5 deg (Ref. 5).

The point on the celestial sphere where the ecliptic crosses the Earth's mean equator (EME) from south to north is called the *ascending node* of the Ecliptic, while the line passing from the Earth's center through that point is called *Earth's vernal equinox* (Υ). The mean Sun is positioned at that point of the sky (as seen from Earth's center) at the time of each spring equinox.

Both the equator and the orbit plane of each planet slowly precess on their respective reference planes, driven by "secular" perturbing effects due to other solar system bodies.

The mean ecliptic regresses, i.e., its nodes slide backwards, on the Invariable Plane at the very slow rate of about 47 seconds of arc per century, while the Earth mean equator regresses on the ecliptic at about 50.26 seconds of arc per annum.

Clearly these motions affect the position of vernal equinox, which is therefore specified as being "of" an epoch, e.g., "of date" (i.e., the present instant of time), or "of 1950.0." The Besselian (tropical) new year has for a long while been considered a convenient defining epoch, usually advanced each half-century, e.g., 1950.0, recently renamed B1950.0, which occurred on January 0.923357, 1950 (i.e., slightly before civil new year 1950, celebrated at 0^h GMT, on January 1.0). The acronym EME50 therefore refers to the Earth mean equator and equinox of B1950.0, while EMO50 specifies the ecliptic plane of that epoch.

Rather than refer to dates frequently quoted in differing calendars, it is customary to specify dates in Julian Days (JD), consecutively counted from an arbitrary date in antiquity (January 1.0, 4713 B.C.). Thus the epoch JD 243 3282.423357 is equivalent to B1950.0.

For each celestial body, a *vernal equinox* may be similarly defined as the ascending node of that body's mean orbit plane (about its primary) upon the body's equator plane. This preferred direction is usually taken to be the X-axis of the inertial coordinate system, centered on that body. The Z-axis coincides with the north-pole axis of the body, and the Y-axis completes a *right-handed* coordinate system.

The equations and related constants defining the above planes are presented in Section IV and may be found in Refs. 4 and 5. Data in Ref. 5 reflect the latest IAU definitions.

A useful way to describe celestial directions is by means of a spherical coordinate system, based either on the equator or else on the mean orbit plane of a body, e.g., that of the Earth or of the target planet. Two angles, declination and right ascension, define the position of any point on the celestial sphere, referred to a planetary equator (Fig. 1).




Fig. 1. Definition of vernal equinox; a position vector R , its declination δ , and right ascension α

Declination (δ) is the central angle formed by the object's position vector with the equator plane, i.e., it is an elevation angle with respect to that plane. It is a form of latitude measure, and is positive towards the north pole and negative to the south of the equator.

Right ascension (α) of an object is the angle in the equator plane between vernal equinox and the projection of the

object's position vector into the equator plane. It is positive in the eastward (i.e., right-handed, counterclockwise, when seen from north) direction, regardless of the planet's (celestial body's) own sense of rotation.

Both of these angles are expressed on the unit sphere. Actual distances between Sun, Earth, and planet can be presented separately and are measured in astronomical units (AU) — a defined mean distance between Sun and Earth (1 AU = 149,597,871 km; see Astrodynamic Constants, Section IV) — or also in kilometers. The distance plot scale in this volume of the handbook is in units of billions of kilometers ($10^9 \times \text{km}$).

Position angles on the celestial sphere expressed with respect to a planetary orbit plane (e.g., the ecliptic) are referred to as latitudes and longitudes.

Ecliptic longitude (L) of an object is the angular distance between Earth's vernal equinox and the projection of the object's position vector into the ecliptic (EMO50), measured in an eastward (i.e., counterclockwise) positive direction.

Sun-Earth-planet (SEP) angle and the corresponding *Earth-Sun-planet (ESP)* angle are defined by the two respective vector pairs: the Earth-Sun and Earth-planet vectors for the SEP-angle, and the Sun-Earth and Sun-planet vectors for the ESP-angle (Fig. 2). The angles are important aids in avoiding periods of communication blackout.




Fig. 2. Definition of Sun- and Earth-to-planet distance, and Earth-Sun- and Sun-Earth-planet angles at any epoch

Cone and clock angles (CA and KA, respectively) are spacecraft-centered coordinates of an object. The polar Z-

axis of the system corresponds to the longitudinal axis of the spacecraft. It is usually pointed either towards the Sun (solar panels must generally be normal to the Sun direction), or to the Earth (the parabolic antenna bore axis should point along the communication link, i.e., towards the Earth). Within this book, the planet-Sun direction is taken to be the Z-axis orientation of the CA, KA system (Fig. 3).




Fig. 3. Definition of cone and clock angles

The prime meridian of this system contains the viewing slot of the spacecraft star tracker, which is usually, but not necessarily, oriented toward the bright southern star Canopus (α -Carinae). Its position on January 1.0, 1950, was at right ascension $\alpha = 95.7103875$ and declination $\delta = -52.667625$ deg, expressed with respect to the EME50. Stellar proper motions slowly change star locations; the method for updating stellar positions as well as data for other stars are provided in the FK4 Star Catalogue (Ref. 6).

ORIGINAL PAGE IS OF POOR QUALITY

The cone angle of an object, described by its spacecraft-centered position vector, is its angular distance from the Z-axis; i.e., it is the Sun-planet-object angle.

The clock angle of an object is measured in the X-Y plane, which is akin to a "spacecraft equator" and normal to the spacecraft Z-axis. Clock angles are measured from the star-tracker (Canopus) meridian; they are positive clockwise when looking along the Z-axis towards the Sun.

Although the clock angle of Canopus is zero by definition (when the spacecraft is Canopus oriented), the Canopus cone angle generally differs from 90 deg, because the position of Canopus is offset several degrees from the ecliptic pole.

Station rise/set times are determined by the daily rotation of the Earth, which takes all ground tracking stations around the polar axis on the small circles that correspond to their terrestrial latitudes. As a result, every object on the celestial sphere rises and sets upon the local horizon of the station (Fig. 4), provided the object's declination is within ± 90 deg of the station latitude (e.g., an object at -55 deg declination cannot be seen by a 40 -deg latitude station). Only objects whose declinations equal the station latitude rise to the zenith, in this daily spectacle (Fig. 5). The angular height of an object above the local horizon is termed *elevation* (Γ). Within this book, an object is considered to have risen if it

reaches above an elevation mask of 6 deg, which accounts for obstructions by local terrain features, atmospheric signal refraction, and radio noise.

Rise and set time information is presented for the Deep Space Network (DSN), the stations for which are listed in Table 1, as well as for three other stations (Table 2).

The Arecibo radio telescope, located within a natural bowl, is not movable. The slewing of a dish, therefore, is replaced by horizontal motion of a cable-mounted cab, which translates the antenna feed to track a celestial object. The resulting daily coverage of this station is clearly very limited.

D. Description of Individual Plots

The individual plots within each annual subsection present the following 26 planetary geometric characteristics as functions of calendar time (recall that for the purposes of this volume "planet" refers to Jupiter):

Plot 1, DECLIN: This plot shows three curves:

- (1) The geocentric declination of the target planet, referred to the Earth mean equator and equinox of B1950.0 (EME50); it is labeled "P" All angular measures on the plots are in degrees.




Fig. 4 Schematic of a distant object's rise and set geometry, with a mask; polar view

ORIGINAL PAGE
OF POOR QUALITY




Fig. 5. Circumpolar and inaccessible regions of the sky for a station observer; meridional view

Table 1. Deep Space Network tracking stations

Name	Geocentric latitude ϕ_s	Geocentric longitude (west) λ_s
DSS 14, Goldstone, California	35.24	116.89
DSS 43, Canberra, Australia	-35.22	211.02
DSS 63, Madrid, Spain	40.24	4.25

Table 2. Non-DSN tracking stations

Name	Geocentric latitude ϕ_s	Geocentric longitude (west) λ_s
Wettzell, Germany	47.69	348.92
Nobeyama, Japan	35.94	221.53
Arecibo, Puerto Rico	18.34	66.75

- (2) The planetocentric (i.e., as seen from the center of the target planet) declination of the Earth, referred to the planet's mean equator and equinox of date (PME/DA). It is labeled "E".
- (3) The planetocentric declination of the Sun, also referred to the PME/DA. It is labeled "S".

Plot II, RT.ASC: Three curves are shown on this plot:

- (4) The geocentric right ascension of the planet with respect to EMES0. It is labeled "P".
- (5) The planetocentric right ascension of the Earth with respect to PME/DA. It is labeled "E".
- (6) The planetocentric right ascension of the Sun with respect to PME/DA. It is labeled "S".

Plot III, DISTANCE: Two curves are presented on this plot:

- (7) The Earth-planet distance, in this volume expressed in units of $10^9 \times \text{km}$ (billions of km). It is labeled "EP".

- (8) The Sun-planet distance, also in billions of km. It is labeled "SP".

Plot IV, EC.LON: A single curve is shown on this plot:

- (9) The planet's heliocentric, ecliptic longitude (with respect to the mean ecliptic and equinox of B1950.0, i.e., EMO50). It is labeled "P".
The annual variation in Earth's heliocentric ecliptic longitude is repetitive to a fair degree of accuracy, hence only a single plot, valid throughout the 1985 to 2020 time period, is provided at the end of the plotted positional data section.

Plot V, SUN-EARTH-PLANET: Two curves are presented:

- (10) The "SEP" angle (Sun-Earth-planet)
(11) The "ESP" angle (Earth-Sun-planet)

Plot VI, CA, KA: Three curves are depicted on the plot:

- (12) The cone angle of Earth for a Sun-oriented spacecraft, labeled "ECA".
(13) The clock angle of Earth, for a Sun-Canopus two-axis-oriented spacecraft. It is labeled "EKA".
(14) The cone angle of Canopus, in any Sun-oriented spacecraft attitude. It is labeled "CCA".

Plot VII, STATION RISE/SET: This plot shows six curves, each representing the Greenwich mean time (GMT) of the following daily Deep Space Network events (all based on a 6-deg mask above the local horizon).

- (15) Planet rise at DSS 14 (Goldstone), labeled "RISE 14".
(16) Planet set at DSS 14, labeled "SET 14".
(17) Planet rise at DSS 43 (Canberra), labeled "RISE 43".
(18) Planet set at DSS 43, labeled "SET 43".
(19) Planet rise at DSS 63 (Madrid), labeled "RISE 63".
(20) Planet set at DSS 63, labeled "SET 63".

Plot VIII, STATION RISE/SET: The plot presents up to six curves, representing the GMT of the following tracking station events (6-deg mask, unless specified otherwise):

- (21) Planet rise at Weilheim, Germany. It is labeled "R WEIL".
(22) Planet set at Weilheim, labeled "S WEIL".
(23) Planet rise at Nobeyama, Japan. It is labeled "R JAPN".

- (24) Planet set at Nobeyama, labeled "S JAPN".
(25) Planet rise at the Arecibo radio telescope, Puerto Rico, above the 70-deg special horizon mask, if realizable. It is labeled "R AREC".
(26) Planet set at Arecibo, under the same conditions as those in (25). It is labeled "S AREC".

III. Application of the Data Presented

The number of practical mission analysis problems to which the information presented in this volume is applicable is obviously large. By no means is it intended to enumerate herein a complete list of such applications. Rather, only some examples, typical of such usage, shall be discussed.

The comments will be arranged in the same order as the graphical output data are presented. Problems in mission analysis to which a particular output type is relevant will be discussed accordingly.

A. Geocentric Declination of Target Body

The spacecraft orbit determination process uses DSN radiometric range and range-rate (Doppler) data. Whenever the observed body as seen from Earth is at near-equatorial (≈ 0 deg) declination, a situation develops that seriously degrades orbit determination accuracy. As a consequence, the timing of critical maneuvers or planetary encounter events should either utilize more sophisticated data types (e.g., Δ DOR, a method based on long-base interferometry) or avoid periods when the target planet is near its node with the Earth's equator (i.e., the times of near-zero declination of the target body).

Communication with the spacecraft greatly depends on a ground station's ability to receive the data signal. An elevation angle high over the horizon, coupled with as long a listening period as possible, are two important assets in this respect. The highest elevation angle of an object at each station occurs at the meridian passage of the object; its value is:

$$\Gamma_{p \max} = 90 - |\phi_s - \delta_p|$$

Its maximum of $\Gamma_p = 90$ deg can be realized only at a station whose latitude, ϕ_s , equals the declination δ_p of the target. Exact equality, however, may cause some data loss: to continue tracking, some antennas may have to swing through a 180-deg arc in azimuth as soon as the object reaches zenith (i.e., at elevation $\Gamma_p = 90$ deg). As already mentioned, the maximum elevation angle drops with an increase in the differ-

ence between station latitude, ϕ_s , and object declination δ_p . At a difference of

$$|\phi_s - \delta_p| + \text{mask} \geq 90 \text{ deg}$$

coverage ceases altogether (Fig. 5). The mask, it will be recalled, is an angle of about 6 deg, designed to assure good signal strength at low elevations.

A notable aspect of this is that, as a station moves in latitude toward either pole (north or south), objects located between the nearer pole and the circumpolar latitude circle δ_c of the station, defined by

$$|\delta_c| = 90 - |\phi_s|$$

will never set and communication may be continuous, if so desired. Knowledge of target-planet declination allows one to choose the particular station for which most critical events would appear at high elevation angles or, conversely, to locate latitudes at which ships and portable or permanent stations should be positioned to support a given difficult mission tracking assignment.

B. Planetocentric Declination of Earth and Sun

These data are based on the target planet's own equator and provide a means of determining the solar lighting conditions at a landing site or during approach. They also allow one to obtain the Earth-oriented downlink elevation angle for a lander on the surface at a given latitude.

For ringed planets (e.g., Saturn), the data can indicate which side (north or south) of the rings is illuminated, which side is visible from Earth, and where on the planetary disk the ring will project its shadow. Observability of disk features and phenomena (e.g., the spokes) may also depend on this information.

C. Right Ascension of Target Planet, Sun, and Earth

This information, when combined with the declination data already discussed, provides positional reference directions toward these bodies, in spherical coordinates as presented, or in Cartesian unit vector form, if transformed as follows:

$$X = \cos \delta_0 \cos \alpha_0$$

$$Y = \cos \delta_0 \sin \alpha_0$$

$$Z = \sin \delta_0$$

where α_0 = right ascension and δ_0 = declination of vector to object.

A number of mission-related problems in spherical trigonometry may be solved using the above information.

D. Sun- or Earth-to-Planet Distance

When the spacecraft is near the planet, Sun-to-planet distance data help in computing such solar effects upon the spacecraft as solar heating and thermal balance, solar-panel electrical output, and spacecraft trajectory and attitude perturbation magnitudes due to the solar-radiation pressure.

Earth-to-planet distance information allows estimation of maximum available data transmission rates in the radio up- or downlink, as well as the one- and two-way light times required in communicating with the spacecraft.

When combined with information on direction unit vectors (X, Y, Z) from paragraph C above, position vectors for Sun, Earth, and planet may be constructed in the respective coordinate systems.

E. Heliocentric Longitude

The heliocentric longitudes of the departure and arrival planetary positions allow sketching the transfer trajectory trace and its orientation with respect to the ecliptic nodes or the perihelion of the target-planet orbit. The relevant planetary information, e.g., the planetary mean orbital elements and related constants, may be found in Section IV.

F. Angles Between Sun, Earth, and Planet (SEP and ESP)

These angles are of paramount importance to spacecraft communication planning. For example, the radio downlink signal becomes distorted and contaminated with noise, if the radio beam passes through the solar corona. This effect is observed near "superior conjunction" between Sun and the target planet. It occurs when the Sun-Earth-planet angle (SEP) reaches near-zero values (commonly, $\leq \pm 5$ deg) while at the same time the Earth-Sun-planet angle (ESP) reaches its maximum value near 180 deg. This unfavorable condition as well as the two other extreme configurations of "opposition" and "inferior conjunction" (neither of which significantly affects up- and downlink communication capability) are shown in Fig. 6.

It is essential to proper mission design that important science, orbit determination, or engineering events be avoided




Fig. 6. Definition of opposition and conjunction events involving Sun, Earth, and planet

during the 10 to 20 days that the spacecraft and/or target planet are near superior conjunction. On the other hand, solar physicists do delight in studying the corona's signature superimposed on the distorted signal. This period of time then becomes an opportunity for corona radio-occultation experiments and related solar studies, including relativity investigations.

G. Cone and Clock Angles

As outlined earlier, the cone- and clock-angle system is primarily useful because it is a spacecraft-fixed set of coordinates. As such, and together with data and equations provided in the Mission Design Handbook series (Ref. 1), the cone and clock angles presented on the plots can be used to resolve a number of geometry problems, such as:

- (1) The angular travel range required by the movable antenna to track the Earth, if Sun-oriented.
- (2) The extent of the Sun-sensor bias required, if the Earth-orientation of the entire spacecraft, together with its antenna, is desired.
- (3) The impact of lighting and thermal constraints upon spacecraft orientation options.

- (4) The Star tracker slot size in cone angle (CA) required to observe Canopus at different times during an orbiter mission.

H. Station Rise and Set

Station rise and set times, given in GMT, greatly affect mission design planning. After consideration is given to the one-way light time required by the radio signal to traverse the Earth-to-planet distance (and back if needed), significant spacecraft events can be positioned in universal (GMT) time using the rise/set information provided, such as to assure that a given station is "up," i.e., that the spacecraft is above mask in elevation angle at a suitable tracking station. The width (in hours) of the up-time band of a station is in itself an indication of the maximum elevation angle available during the station's pass "under" the spacecraft (i.e., the planet) — the greater the width, the higher the maximum elevation angle.

IV. Astrodynamic Constants

This section presents information necessary for relating the various coordinate systems used in the handbook. For other astrodynamic constants, see Refs. 1, 4, 5, and 7.


ORIGINAL PAGE IS OF POOR QUALITY

Mean orbital elements for the planets are *averaged* analytical models of planetary motion, adopted for reference usage by the astronomical community (e.g., H. M. Nautical Almanac Office or the U.S. Naval Observatory). Mean elements change as functions of time, T_{50} .

T_{50} is defined as the time interval, in Julian centuries, elapsed at date of interest, since epoch January 1.0, 1950, 0^h ephemeris time (ET) (i.e., Julian epoch J1950.0 = JD2433282.5). A Julian century contains 36525.0 mean solar (i.e., calendar) days. Ephemeris time is an atomically controlled uniform time, approximated as: ET = GMT + DUT, where the time increment DUT = +54.2 s (as of July 1983). DUT is currently increasing at about one second per year.

Six elements completely define the mean orbit (Fig. 7), as follows (all angular measures in degrees):

(a) CELESTIAL SPHERE VIEW



(b) ORBIT PLANE VIEW




Fig. 7. Definition of the mean elements of a planetary orbit

a = semimajor axis (in astronomical units, 1 AU = 149,597,871 km).

e = eccentricity of orbit (dimensionless).

M = mean anomaly, a mean motion angular position from perihelion (i.e., closest approach to Sun) at time T_{50} (for numerical convenience also expressed as d_{50} , days from January 1.0, 1950 0^h ET). True anomaly (actual position angle from perihelion) can be obtained from M by way of iteration using Kepler's transcendental equation.

i_{50} = inclination angle of mean orbit plane at time T_{50} , with respect to the mean ecliptic of epoch B1950.0 (EMO50).

Ω_{50} = longitude of ascending node of mean orbit upon the ecliptic of B1950.0, measured from Earth's vernal equinox.

ω_{50} = argument of perihelion, an angle measured from the ascending node of the mean orbit on the ecliptic of B1950.0 to the perihelion of the mean orbit of the planet.

As presented by F. Sturms (Ref. 4) and based on information in the Explanatory Supplement to the Ephemeris (Ref. 7), the mean elements are:

(1) Earth:

$$a = 1.00000023 \text{ AU} = 149,597,905 \text{ km}$$

$$e = 0.0167301085 - 0.000041926 T_{50} - 0.000000126 T_{50}^2$$

$$M = 358.000682 + 0.9856002628 d_{50} - 0.0001550000 T_{50}^2 - 0.0000033333 T_{50}^3$$

$$i_{50} = 0.013076 T_{50} - 0.000009 T_{50}^2$$

$$\Omega_{50} = 174.40956 - 0.24166 T_{50} + 0.00006 T_{50}^2$$

$$\omega_{50} = 287.67097 + 0.56494 T_{50} + 0.00009 T_{50}^2$$

(2) Jupiter:

$$a = 5.202803 \text{ AU} = 778,328,250 \text{ km}$$

$$e = 0.04841911 + 0.00016302 T_{50}$$

$$M = 302.650461 + 0.0830898769 d_{50}$$

$$i_{50} = 1.30592 - 0.00205 T_{50} + 0.00003 T_{50}^2$$

$$\Omega_{50} = 99.94335 + 0.16728 T_{50} + 0.00055 T_{50}^2$$

$$\omega_{50} = 273.57374 + 0.04756 T_{50} - 0.00086 T_{50}^2$$

ORIGINAL PAGE IS
OF POOR QUALITY

The equatorial plane of a celestial body at an epoch of interest, T_{50} , was defined by the International Astronomical Union Working Group on Cartographic Coordinates and Rotational Elements (Ref. 5) by the right ascension α_{50} and declination δ_{50} of the body's "north" polar axis direction with respect to the Earth mean equator of epoch B1950.0 (EME50), as shown in Fig. 8. The position of the rotating prime meridian of the body is defined by an angle W , measured positive counterclockwise in that body's equatorial plane from the ascend-

ing node of that equator upon the EME50, to the prime meridian position at epoch d_{50} . According to the Ref. 5, Table I:

(1) Sun pole:

$$\begin{aligned}\alpha_{50} &= 285.90 \text{ (deg)} \\ \delta_{50} &= 63.90 \\ W &= 240.90 + 14.1844000 d_{50}\end{aligned}$$

(2) Earth pole:

$$\begin{aligned}\alpha_{50} &= 0.00 - 0.640 T_{50} \\ \delta_{50} &= 90.00 - 0.557 T_{50} \\ W &= 99.87 + 360.9856123 d_{50}\end{aligned}$$

(3) Jupiter pole:

$$\begin{aligned}\alpha_{50} &= 268.00 - 0.008 T_{50} \\ \delta_{50} &= 64.50 + 0.003 T_{50} \\ W &= 80.60 + 870.5360000 d_{50} \text{ (System III, radio longitude)}\end{aligned}$$

(4) Invariable Plane pole:

$$\begin{aligned}\alpha_{50} &= 272.40 \\ \delta_{50} &= +66.99\end{aligned}$$

Note that in the computation of W , care should be taken not to lose significant digits during arithmetic calculations.

Positional data in this handbook were obtained using the JPL DE-118 precision planetary ephemeris on magnetic tape in tabular format (Ref. 3).




Fig. 8. IAU definition of planetary pole and prime meridian at time d_{50} , elapsed since January 1.0, 1950 0^h ET

Acknowledgments

The contributions, reviews, and suggestions by members of the Handbook Advisory Committee, especially those of K. T. Nock, R. E. Diehl, W. I. McLaughlin, W. E. Bollman, P. A. Penzo, R. A. Wallace, D. F. Bender, D. V. Byrnes, L. A. D'Amario, T. H. Sweetser, and R. S. Schlaifer are acknowledged and greatly appreciated. The authors would like to thank Mary Fran Buehler and David E. Fulton for their editorial contribution.

References

1. Sergeyevsky, A. B., et al., *Interplanetary Mission Design Handbook*, JPL Publication 82-43, Volume 1, Parts 1-4. Jet Propulsion Laboratory, Pasadena, Calif., March 1, 1983.
2. Sergeyevsky, A. B., *Mission Design Data for Venus, Mars, and Jupiter through 1990*, Technica. Memorandum 33-736. Jet Propulsion Laboratory, Pasadena, Calif., Sept. 1, 1975.
3. Newhall, XX, Standish, E. M., and Williams, J. G., "DE-102: A Numerically Integrated Ephemeris of the Moon and Planets, Spanning 44 Centuries," *Astronomy and Astrophysics*, 1983 (in press).
4. Sturms, F. M., *Polynomial Expressions for Planetary Equators and Orbit Elements With Respect to the Mean 1950.0 Coordinate System*, Technical Report 32-1508, pp. 6-9. Jet Propulsion Laboratory, Pasadena, Calif., Jan. 15, 1971.
5. Davies, M. E., et al., "Report of the IAU Working Group on Cartographic Coordinates and Rotational Elements of the Planets and Satellites: 1982," *Celestial Mechanics*, Vol. 29, No. 4, pp. 309-321, April 1983.
6. Fricke, W., et al., *Fourth Fundamental Catalogue (FK 4)*, No. 10 in the series, *Veröffentlichungen des Astronomischen Rechen-Instituts Heidelberg*, Verlag L. Braun, Karlsruhe, 1963.
7. "Explanatory Supplement to the Ephemeris", Her Majesty's Stationery Office, London, 1961.

Positional Data

Jupiter
1985—2020

PRECEDING PAGE BLANK NOT FILMED

PAGE 12 . INTENTIONALLY BLANK

Jupiter

1985



PRECEDING PAGE BLANK NOT FILMED

14 INTENTIONALLY BLANK

DECLIN RT.ASC 1985

ORIGINAL PAGE IS
OF POOR QUALITY



JUPITER 1985



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1985



DISTANCE
EC.LON
1985



SEP, ESP
CA, KA
1985

JUPITER 1985


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1985

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 1985



STA R/S
NON-DSN
1985

ORIGINAL PAGE 1
OF POOR QUALITY

JUPITER 1985





Jupiter

1986

DECLIN RT.ASC 1986

JUPITER 1986



ORIGINAL PREDICTION
OF POOR QUALITY



ORIGINAL PAGE 73
OF POOR QUALITY

JUPITER 1986

DISTANCE
EC.LON
1986





SEP, ESP
CA, KA
1986

JUPITER

1986


ORIGINAL PAGE
OF POOR QUALITY



STA R/S
DSN
1986

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 1986



STA R/S
NON-DSN
1986

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1986



Jupiter



1987

DECLIN RT.ASC 1987

JUPITER

1987



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 19
OF POOR QUALITY

JUPITER 1987



DISTANCE
EC.LON
1987



**SEP, ESP
CA, KA
1987**

JUPITER 1987


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1987

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 1987



STA R/S
NON-DSN
1987

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1987



Jupiter



1988

DECLIN
RT.ASC
1988

JUPITER

1988

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER

1988

DISTANCE
EC.LON
1988





SEP, ESP
CA, KA
1988

JUPITER

1988


ORIGINAL PAGE 10
OF POOR QUALITY



STA R/S
DSN
1988

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 1988



STA R/S
NON-DSN
1988

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1988



Jupiter



1989

DECLIN RT.ASC 1989

JUPITER

1989



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 1989

DISTANCE
EC.LON
1989





SEP, ESP
CA, KA
1989

JUPITER

1989


ORIGINAL PAGE 13
OF POOR QUALITY



STA R/S
DSN
1989

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 1989



ORIGINAL PAGE IS
OF POOR QUALITY

The graph displays the station rise/set times in GMT for five stations over a period from January 1, 1989, to January 16, 1990. The Y-axis represents the time in GMT hours, ranging from 0 to 24. The X-axis represents the date in YR/M/D format. Five lines are plotted, each representing a different station. The lines show a general downward trend, indicating that the station rise/set times are decreasing over the period shown. The stations are labeled as follows: AREC, JAPN, WEIL, and two unlabeled stations.

Station	89/01/01	89/01/21	89/02/10	89/03/02	89/03/22	89/04/11	89/05/01	89/05/21	89/06/10	89/06/30	89/07/20	89/08/09	89/08/29	89/09/18	89/10/08	89/10/28	89/11/17	89/12/07	89/12/27	90/01/16
AREC	24.0	22.0	20.0	18.0	16.0	14.0	12.0	10.0	8.0	6.0	4.0	2.0	0.0							
JAPN	18.0	16.0	14.0	12.0	10.0	8.0	6.0	4.0	2.0	0.0										
WEIL	13.5	11.5	9.5	7.5	5.5	3.5	1.5	0.0												
Unlabeled 1	24.0	22.0	20.0	18.0	16.0	14.0	12.0	10.0	8.0	6.0	4.0	2.0	0.0							
Unlabeled 2	24.0	22.0	20.0	18.0	16.0	14.0	12.0	10.0	8.0	6.0	4.0	2.0	0.0							



Jupiter

1990

DECLIN RT.ASC 1990

JUPITER 1990



ORIGINAL PAGE 13
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 1990



DISTANCE
EC.LON
1990



SEP, ESP
CA, KA
1990

JUPITER 1990


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1990

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 1990



STA R/S
NON-DSN
1990

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 1990



Jupiter



1991

DECLIN
RT.ASC
1991

JUPITER



1991

ORIGINAL PAGE 19
OF POOR QUALITY



JUPITER



1991

ORIGINAL PAGE IS
OF POOR QUALITY**DISTANCE**
EC.LON
1991

SEP, ESP
CA, KA
1991

JUPITER 1991


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1991

ORIGINAL PAGE 17
OF POOR QUALITY


JUPITER 1991



STA R/S
NON-DSN
1991

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1991





Jupiter

1992

DECLIN RT.ASC 1992

JUPITER 1992



ORIGINAL PAGE 13
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 1992



DISTANCE
EC.LON
1992



SEP, ESP
CA, KA
1992

JUPITER 1992


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1992

ORIGINAL PALETTE
OF POOR QUALITY


JUPITER 1992



**STA R/S
NON-DSN
1992**

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1992





Jupiter

1993

DECLIN RT.ASC 1993

JUPITER 1993



ORIGINAL PAGE 13
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1993



**DISTANCE
EC.LON
1993**



SEP, ESP
CA, KA
1993

JUPITER 1993


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1993

ORIGINAL PAGE 11
OF POOR QUALITY


JUPITER 1993



STA R/S
NON-DSN
1993

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1993





Jupiter

1994

DECLIN RT.ASC 1994



JUPITER 1994

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS JUPITER 1994
OF POOR QUALITY



DISTANCE EC.LON 1994



SEP, ESP
CA, KA
1994

JUPITER 1994


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1394

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 1994



STA R/S
NON-DSN
1994

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 1994





Jupiter

1995

DECLIN
RT.ASC
1995

JUPITER 1995



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1995



DISTANCE
EC.LON
1995



SEP, ESP
CA, KA
1995

JUPITER 1995


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1995

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 1995



STA R/S
NON-DSN
1995

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1995



Jupiter



1996

DECLIN RT.ASC 1996

JUPITER



1996

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY JUPITER 1996



DISTANCE
EC.LON
1996



SEP, ESP
CA, KA
1996

JUPITER 1996


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
1996


JUPITER 1996



STA R/S
NON-DSN
1996

ORIGINAL PAGE 19
OF POOR QUALITY

JUPITER 1996



Jupiter



1997

DECLIN RT.ASC 1997

JUPITER

1997



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1997



DISTANCE
EC.LON
1997



SEP, ESP
CA, KA
1997

JUPITER 1997


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1997

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 1997



**STA R/S
NON-DSN
1997**

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 1997





Jupiter

1998

DECLIN
RT.ASC
1998

JUPITER 1998



ORIGINAL PAGE 18
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1998

DISTANCE
EC.LON
1998





SEP, ESP
CA, KA
1998

JUPITER

1998


ORIGINAL PAGE 15
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

STA R/S
DSN
1998


JUPITER 1998



STA R/S
JON-DSN
1998

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1998



Jupiter



1999

DECLIN RT.ASC 1999

JUPITER

1999



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1999



DISTANCE
EC.LON
1999



SEP, ESP
CA, KA
1999

JUPITER 1999


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
1999

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 1999



STA R/S
NON-DSN
1999

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 1999





Jupiter

2000

DECLIN RT.ASC 2000

JUPITER 2000



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2000



DISTANCE
EC.LON
2000



SEP, ESP
CA, KA
2000

JUPITER 2000


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2000

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 2000



**STA R/S
NON-DSN
2000**

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2000





Jupiter

2001

DECLIN RT.ASC 2001

JUPITER 2001



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2001



DISTANCE
EC.LON
2001



SEP, ESP
CA, KA
2001

JUPITER 2001


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2001

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2001



STA R/S
NON-DSN
2001

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2001





Jupiter

2002

DECLIN RT.ASC 2002

JUPITER 2002



ORIGINAL PAGE 13
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2002



DISTANCE
EC.LON
2002



SEP, ESP
CA, KA
2002


JUPITER 2002

ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2002



STA R/S
NON-DSN
2002

ORIGINAL PAGE 10
OF POOR QUALITY

JUPITER 2002





Jupiter



2003

DECLIN RT.ASC 2003

JUPITER 2003

ORIGINAL PAGE IS
OF POOR QUALITY







SEP, ESP
CA, KA
2003

JUPITER

2003


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 12
OF POOR QUALITY

STA R/S
DSN
2003


JUPITER 2003



STA R/S
NON-DSN
2003

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2003





Jupiter

2004

DECLIN RT.ASC 2004

JUPITER 2004



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2004



DISTANCE
EC.LON
2004



SEP, ESP
CA, KA
2004

JUPITER 2004


ORIGINAL PAGE 1
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

STA R/S
DSN
2004


JUPITER 2004



**STA R/S
NON-DSN
2004**

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2004



Jupiter



2005

DECLIN RT.ASC 2005

JUPITER

2005

ORIGINAL PAGE IS
OF POOR QUALITY





ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER

2005

DISTANCE
EC.LON
2005





SEP, ESP
CA, KA
2005

JUPITER

2005


ORIGINAL PAGE 13
OF POOR QUALITY



STA R/S
DSN
2005

ORIGINAL PICTURE
OF POOR QUALITY


JUPITER 2005



STA R/S
NON-DSN
2005

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2005





Jupiter

2006

DECLIN
RT.ASC
2006

JUPITER 2006



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2006

DISTANCE
EC.LON
2006





SEP, ESP
CA, KA
2006

JUPITER

2006


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
2006


JUPITER 2006



STA R/S
NON-DSN
2006

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2006





Jupiter

2007

DECLIN RT.ASC 2007

JUPITER 2007



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2007

DISTANCE
EC.LON
2007





SEP, ESP
CA, KA
2007

JUPITER

2007


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
2007


JUPITER 2007



STA R/S
NON-DSN
2007

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2007



Jupiter



2008

DECLIN
RT.ASC
2008

JUPITER

2008



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 19
OF POOR QUALITY

JUPITER 2008

DISTANCE
EC.LON
2008





SEP, ESP
CA, KA
2008

JUPITER

2008


ORIGINAL PAGE 13
OF POOR QUALITY



STA R/S
DSN
2008

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 2008



STA R/S
NON-DSN
2008

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2008



Jupiter



2009



DECLIN RT.ASC 2009

JUPITER

2009

ORIGINAL PAGE 13
OF POOR QUALITY







**SEP, ESP
CA, KA
2009**

JUPITER 2009


ORIGINAL PAGE 10
OF POOR QUALITY



STA R/S
DSN
2009

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2009



STA R/S
NON-DSN
2009

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2009





Jupiter

2010

DECLIN
RT.ASC
2010

JUPITER 2010



ORIGINAL PAGE 19
OF POOR QUALITY



ORIGINAL PAGE 10
OF POOR QUALITY

JUPITER 2010



DISTANCE
EC.LON
2010



SEP, ESP
CA, KA
2010

JUPITER 2010


ORIGINAL PAGE 13
OF POOR QUALITY



STA R/S
DSN
2010

ORIGINAL PAGE 18
OF POOR QUALITY


JUPITER 2010



**STA R/S
NON-DSN
2010**

ORIGINAL FILED
OF POOR QUALITY

JUPITER 2010





Jupiter



2011

DECLIN RT.ASC 2011

JUPITER 2011

ORIGINAL PAGE 13
OF POOR QUALITY







SEP, ESP
CA, KA
2011

JUPITER 2011


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2011

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 2011



STA R/S
NON-DSN
2011

ORIGINAL PAGE 19
OF POOR QUALITY

JUPITER 2011



Jupiter



2012

DECLIN RT.ASC 2012

JUPITER

2012



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2012

DISTANCE
EC.LON
2012





SEP, ESP
CA, KA
2012

JUPITER

2012


ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE IS
OF POOR QUALITY

STA R/S
DSN
2012


JUPITER 2012



STA R/S
NON-DSN
2012

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2012



Jupiter



2013

DECLIN RT.ASC 2013

JUPITER

2013



ORIGINAL PAGE 15
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2013



DISTANCE
EC.LON
2013



SEP, ESP
CA, KA
2013

JUPITER 2013


ORIGINAL PAGE 12
OF POOR QUALITY



STA R/S
DSN
2013

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2013



STA R/S
NON-DSN
2013

ORIGINAL PAGE 12
OF POOR QUALITY

JUPITER 2013



Jupiter



2014

C - 3

DECLIN RT.ASC 2014

JUPITER 2014



ORIGINAL PAGE
OF POOR QUALITY



ORIGINAL PRICE IN
OF POOR QUALITY

JUPITER 2014

DISTANCE
EC.LON
2014





SEP, ESP
CA, KA
2014

JUPITER

2014


ORIGINAL PAGE 11
OF POOR QUALITY



STA R/S
DSN
2014

ORIGINAL DRAWING
OF POOR QUALITY


JUPITER 2014



STA R/S
NON-DSN
2014

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2014





Jupiter

2015

DECLIN RT.ASC 2015

JUPITER 2015

ORIGINAL PAGE 13
OF POOR QUALITY





ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER

2015

DISTANCE
EC.LON
2015





SEP, ESP
CA, KA
2015

JUPITER

2015


ORIGINAL PAGE 15
OF POOR QUALITY



STA R/S
DSN
2015

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 2015



STA R/S
NON-DSN
2015

ORIGINAL PAGE 15
OF POOR QUALITY

JUPITER 2015



Jupiter



2016

DECLIN RT.ASC 2016



JUPITER

2016

ORIGINAL PAGE IS
OF POOR QUALITY



DISTANCE EC.LON 2016





SEP, ESP
CA, KA
2016

JUPITER

2016


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2016

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2016



STA R/S
NON-DSN
2016

ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2016





Jupiter

2017

DECLIN RT.ASC 2017

JUPITER 2017



ORIGINAL PAGE 10
OF POOR QUALITY



ORIGINAL PAGE 13
OF POOR QUALITY

JUPITER 2017



DISTANCE
EC.LON
2017



SEP, ESP
CA, KA
2017

JUPITER 2017


ORIGINAL PAGE 13
OF POOR QUALITY



STA R/S
DSN
2017

ORIGINAL PAGE 13
OF POOR QUALITY


JUPITER 2017



STA R/S
NON-DSN
2017

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2017



Jupiter



2018

DECLIN
RT.ASC
2018

JUPITER

2018



ORIGINAL PAGE 13
OF POOR QUALITY



JUPITER

2018 ORIGINAL PAGE IS
OF POOR QUALITY



DISTANCE
EC.LON
2018



SEP, ESP
CA, KA
2018

JUPITER 2018


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2018

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2018



STA R/S
NON-DSN
2018

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2018





Jupiter

2019

**DECLIN
RT.ASC
2019**

JUPITER 2019



ORIGINAL PAGE IS
OF POOR QUALITY



ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2019



DISTANCE
EC.LON
2019



SEP, ESP
CA, KA
2019

JUPITER 2019


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2019

ORIGINAL PAGE IS
OF POOR QUALITY


JUPITER 2019



STA R/S
NON-DSN
2019

ORIGINAL PAGE IS
OF POOR QUALITY

JUPITER 2019



Jupiter



2020

DECLIN RT.ASC 2020

JUPITER



2020

ORIGINAL PAGE IS
OF POOR QUALITY



JUPITER



2020

ORIGINAL PAGE IS
OF POOR QUALITYDISTANCE
EC.LON
2020

SEP, ESP
CA, KA
2020

JUPITER 2020


ORIGINAL PAGE IS
OF POOR QUALITY



STA R/S
DSN
2020

ORIGINAL PAGE 19
OF POOR QUALITY


JUPITER 2020



**STA R/S
NON-DSN
2020**

ORIGINAL PAGE 18
OF POOR QUALITY

JUPITER 2020




Positional Data

**Earth
1985—2020**

EC.LON EARTH ALL YEARS

ORIGINAL PAGE IS
OF POOR QUALITY



PRECEDING PAGE BLANK NOT FILMED

PAGE 232

PALLY BLANK